



RESEARCH

Open Access

How echinococcosis affects potential cancer markers in plasma: galectin-3, sN-cadherin and sE-cadherin? A preliminary report

Joanna Giebultowicz^{1*}, Malgorzata Polanska-Plachta², Piotr Wroczynski¹, Piotr Zaborowski³ and Jerzy A Polanski²

Abstract

Background: An increasing number of publications are suggesting that galectin-3 (Gal-3) and soluble cadherin fragments, such as E-cadherin (sE-CAD) and N-cadherin (sN-CAD), may be considered as cancer markers. Despite the promising results of the studies, there are no data concerning their levels in the plasma of echinococcosis patients. In most cases, echinococcosis affects the liver, and its symptoms and disease course are very similar to those of liver cancer. The aim of the present study was to observe whether echinococcosis affects the concentration of soluble sN-CAD, sE-CAD fragments and Gal-3 in plasma and to determine which of them could be considered reliable liver cancer markers for further research.

Methods: The concentrations of sN-CAD, sE-CAD and Gal-3 in the EDTA plasma of patients suffering from echinococcosis ($N = 20$), liver cancers ($N = 10$) and healthy subjects ($N = 20$) were measured using the ELISA method.

Results: The plasma concentration of sE-CAD was lower ($p = 0.0381$), and that of Gal-3 higher ($p = 0.0288$), in echinococcosis than in the healthy group. However, only the concentration of sE-CAD differed significantly among the three analysed groups. In echinococcosis there was a correlation between the sE-CAD and CRP levels ($r_s = 0.79$; $p = 0.0066$) as well as a correlation between the sE-CAD level and the number of leukocytes ($r_s = 0.65$; $p = 0.0210$) in the blood.

Conclusions: Echinococcosis affects the concentration of soluble sE-CAD fragments and Gal-3 in plasma. sE-CAD can be considered as a marker for differentiation between liver cancer and echinococcosis, a parasitic liver disease similar in symptoms. Further study is required to confirm these preliminary results.

Virtual Slides: The virtual slide(s) for this article can be found here:
<http://www.diagnosticpathology.diagnomx.eu/vs/2115657402650448>.

Keywords: Echinococcosis, N-cadherin, E-cadherin, Galectin-3, Liver cancer

Background

Echinococcosis is a parasitic disease caused by a cestode belonging to the genus *Echinococcus*. The most important species for medical purposes are *E. granulosus* (cystic echinococcosis) and *E. multilocularis* (alveolar echinococcosis). The liver is the most commonly invaded organ (the liver and the lungs represent more than 90% of reported cases). Pathological symptoms such as

abdominal pain and/or jaundice are caused by the pressure the enlarged cyst exerts on the liver tissue. There is additionally a palpable mass in the hepatic area [1]. Similar symptoms are registered in patients with liver cancer. Moreover, echinococcosis and liver cancer have a long asymptomatic occurrence. Cystic echinococcosis is usually easy to diagnose using an ultrasound image. It is, however, occasionally difficult to distinguish it from other liver tumours. In these cases, other diagnostic tools, such as serological tests, are used. False positive or false negative reactions, however, may still occur [2]. Moreover, echinococcosis is sometimes not considered

* Correspondence: joanna.giebultowicz@wum.edu.pl

¹Department of Bioanalysis and Drugs Analysis, Faculty of Pharmacy, Medical University of Warsaw, 1 Banacha Street, Warsaw PL-02097, Poland
Full list of author information is available at the end of the article

during medical diagnosis due to the incidence rates of the two diseases. An erroneous diagnosis of a cyst as being a cancer is associated with the risk of its being damaged during surgery. This can result in anaphylactic shock. Proper cancer markers should therefore be able to distinguish between the cancer and the parasitic cyst.

An increasing number of publications are suggesting that galectin-3 (Gal-3) and soluble cadherin fragments, such as E-cadherin (sE-CAD) and N-cadherin (sN-CAD), may be considered as cancer markers. Although the concentration of proper cancer markers should not only distinguish between healthy and cancerous subjects, but also between cancerous subjects and those suffering from other tumour progression diseases such as echinococcosis, there are no data concerning sE-CAD, sN-CAD and Gal-3 levels in the plasma of echinococcosis patients. This disease may sometimes be misdiagnosed as cancer, which can result in patient death.

The aim of the present study was to observe whether echinococcosis affects the concentration of soluble sN-CAD, sE-CAD fragments and Gal-3 in plasma and to determine which of them are suitable for further research as reliable liver cancer markers.

Methods

Materials

Mouse anti-N-cadherin antibodies and 3,3',5,5'-Tetramethylbenzidine (TMB) solution for ELISA were purchased from Sigma-Aldrich (Poznan, Poland). Goat anti-mouse antibodies conjugated with horseradish peroxidase were supplied by Jackson ImmunoResearch Laboratories (Biokom, Janki, Poland). Recombinant Human N-Cadherin/Fc Chimera and Quantikine Human sE-Cadherin Immunoassay were obtained from R&D Systems (Biokom, Janki, Poland). The Gal-3 ELISA kit was purchased from BioVendor R&D (Biokom, Janki, Poland) and the hs-CRP kit from Demeditec Diagnostic (Biokom, Janki, Poland).

Ethics

This study was approved by the Polish Research Ethics Committee (KB 32/2011).

Apparatus

Photometric measurements were performed using a Bio-Tek Synergy 5 spectrophotometer (Biokom, Janki, Poland).

Plasma collection

EDTA-plasma samples were obtained from healthy volunteers ($n = 20$) and from subjects suffering from cystic echinococcosis ($n = 20$) as well as primary (hepatocellular carcinoma, cholangiocarcinoma) and

secondary (metastatic adenocarcinoma of the colon and metastatic endometrial cancer) liver cancer ($n = 10$), treated at the Warsaw Hospital for Infectious Diseases and at the Czerniakowski Hospital, Medical University of Warsaw. Plasma samples were separated from blood within 1 to 3 h after blood collection and separated by centrifugation at 2500 rpm for 30 min at about 10°C. The separated samples were stored at -70°C until being assayed.

ELISA methods

The concentration of sE-CAD, Gal-3 and hs-CRP in plasma was determined by sandwich ELISA using commercially available kits. As no commercially available kit to measure plasma sN-CAD concentration was available, measurements were performed using the procedure reported by Derycke for direct ELISA [3]. The ELISA plates were coated with twice-diluted plasma overnight. Mouse monoclonal anti-N-cadherin antibodies (1:200) were the primary antibody and goat anti-mouse antibodies conjugated with horseradish peroxidase (1:5000) were the secondary antibody. TMB was used as a substrate. The absorbance was recorded at 450 nm, with the correction wavelength at 620 nm, after the reaction was quenched with an acidic solution (2 M aq. HCl).

Statistical analysis

Normal distribution and variance homogeneity of the data (before and after the variables had been transformed into their natural logarithms) were checked using the Shapiro-Wilk test and Leven's test, respectively. When the data were not normally distributed or their variance not homogeneous ($p < 0.05$), the statistical significance evaluation was performed using the Kruskal-Wallis test. Otherwise, the ANOVA test was used. The correlations between the variables were analyzed using Spearman's rank correlation test, and expressed as r_s (correlation coefficient) and p ($p < 0.05$ were described as significant). Discriminant function analysis was used to determine which variables discriminate between the examined groups. The classification functions were used to predict a categorical dependent variable by one or more variables. Each function gives the formula according to which the classification scores for the analyzed case can be calculated as follows:

$$Si = ci + wi1 \times x1 + wi2 \times x2 + \dots + wim \times xm$$

where: si - is the score for i 'th group of 1,2. . . m variables; ci - constant for the i 'th group; $wi1 \dots wim$ - discriminant function coefficients.

Each case is classified into the group with the highest score. Calculations were performed using STATISTICA version 9.1 software.

Results

Analyzed group features

The mean CRP level was 5.0 (7.3) mg/dl in the cancerous patients and 4.9 (5.5) mg/dl ($p = 0.59$; NS) in those suffering from echinococcosis. Their leucocytes levels were 10.5 (3.8) and 5.3 (1.6) K/ μ l ($p = 0.0022$), respectively. The mean age was 64 yrs (range 38-79) in the cancerous group, 48 yrs (range 21-75) in the echinococcosis group, and 41 yrs (range 28-65) in the control group. Occurrence of Echinococcosis was confirmed using ultrasound imaging and serological tests. Cancers were confirmed histopathologically.

sN-CAD concentration

Average plasma concentration of sN-CAD in the examined groups is summarized in Table 1. Post-hoc testing revealed that the sN-CAD concentration in the plasma of healthy subjects was significantly higher than in the plasma of echinococcosis ($p = 0.0019$) and liver cancer ($p < 0.00001$) patients. The difference between the sN-CAD level in echinococcosis and liver cancer was not significant ($p = 0.26$). There was also no statistically significant correlation between sN-CAD and CRP levels or between the sN-CAD level and the number of leukocytes in the examined groups.

sE-CAD concentration

Average plasma concentration of sE-CAD in the examined groups is summarized in Table 1. Post-hoc testing revealed that the sE-CAD concentration in the plasma of cancerous subjects was significantly higher than in the plasma of echinococcosis ($p = 0.0001$) patients and healthy subjects ($p = 0.0381$). Moreover, in echinococcosis, there was a correlation between the sE-CAD and CRP levels ($r_s = 0.79$; $p = 0.0066$) as well as a correlation between the sE-CAD level and the number of leukocytes ($r_s = 0.65$; $p = 0.0210$) in the blood.

Gal-3 concentration

Average plasma concentration of Gal-3 in the examined groups is summarized in Table 1. Post-hoc NIR testing revealed that the Gal-3 concentration in the plasma of echinococcosis patients was higher than that in the plasma of healthy subjects ($p = 0.0211$) and cancerous

patients ($p = 0.1151$, NS). There were no significant differences between the concentration of Gal-3 in the plasma of cancerous and healthy subjects ($p = 0.65$, NS). There was also no statistically significant correlation between Gal-3 and CRP levels or between the Gal-3 level and the number of leukocytes in the examined groups.

Discriminant analysis

Discriminant function analysis showed that by involving data such as the plasma concentration of Gal-3 and sN-CAD, 100% of healthy cases, 71% of echinococcosis cases, and 67% of liver cancer cases could be correctly classified.

The test of dimensionality for discriminant analysis indicated that only the first dimension was statistically significant ($p = 0.00001$). The standardized canonical discriminant function coefficients were -1.06 for sN-CAD and 0.58 for Gal-3 plasma concentration. The mean canonical roots were -1.44 for the healthy subjects, 0.31 for the echinococcosis patients and 1.13 for the cancer patients. The classification function coefficients are summarized in Table 2.

Discussion

Galectin-3 is reported to be overexpressed in a variety of neoplastic cells and is suggested to be involved in tumour metastasis by, e.g. enhancing adhesion between tumour cells and extracellular matrix, promoting embolization. However, no generalized conclusion on the role of Gal-3 in cancer has yet been established. The lectin is not only produced and secreted into the plasma by tumour tissues, but also by peritumoural inflammatory cells and stromal cells.

Elevated Gal-3 levels in the serum/plasma have been reported in some cancers, e.g., metastatic breast and gastrointestinal cancer. However, decreased levels of this lectin have been observed in metastatic breast, endometrial and ovary carcinomas, while the levels in Non-Hodgkin's lymphoma and melanoma [4] have not been observed to change. Elevated levels of circulating Gal-3 have not only been observed in cancers, but also in chronic inflammations, e.g. inflammatory bowel disease [5], obesity, type 2 diabetes [6], and heart failure [7,8].

Table 1 Average values (mean, median) and dispersion (standard deviation, interquartile range) of soluble N-cadherin, E-cadherin and galectin-3 concentration in the plasma of the examined group

Concentration [ng/ml]	Echinococcosis (n = 20)		Cancer (n = 10)		Healthy (n = 20)	
	Mean (SD)	Median (IQR)	Mean (SD)	Median (IQR)	Mean (SD)	Median (IQR)
sN-CAD	94 (40) * ^H	80 (23)	58 (47) * ^H	50 (70)	141 (11) * ^{EC}	146 (12)
sE-CAD	9 (11)	6.5 (3.2) * ^{CH}	12.4 (4.6)	11.0 (1.5) * ^{EH}	8.6 (1.4)	8.4 (2.0) * ^{EC}
Gal-3	12.5 (6.0) * ^{CH}	10.4 (8.8)	9.9 (2.7) * ^E	9.1 (4.1)	9.1 (1.5) * ^E	9.8 (2.3)

*- statistically significant differences ($p < 0.05$) between the examined groups (E-echinococcosis, C-cancer, H-healthy)

Table 2 Classification function coefficients determined using discriminant function analysis

Variable	Echinococcosis	Cancer	Healthy
sN-CAD [ng/ml]	0.045	0.020	0.093
Gal-3 [ng/ml]	0.550	0.488	0.196
Constant	-6.439	-4.292	-8.709

The present studies showed that a higher circulating Gal-3 concentration was observed in liver cancer (NS) and echinococcosis patients than in the healthy control group. The results concerning the cancerous patients, even if not significant, seem to be consistent with the results mentioned above [4] as well as with those reported by Matsuda et al. [9], who observed a higher level of the circulating lectin in liver cancer than in the control group and subjects with chronic liver disease. However, the present report would seem to be the first to compare the levels of Gal-3 in the plasma of cancer patients, echinococcosis patients and healthy volunteers. The highest level observed in echinococcosis could be the result of inflammation related to parasite invasion. Nevertheless, liver cancer is also associated with inflammation and exhibited similar CRP levels to, and higher leukocyte levels than, echinococcosis. Factors other than the inflammation that occurs in echinococcosis might therefore influence the Gal-3 level in the plasma.

sE-cadherin is a form, cleaved from E-cadherin by proteases such as MMP-3, MMP-7 [10], plasmin [11], ADAM10 [12], ADAM15 [13], whereas sN-CAD is cleaved by ADAM10 [14], MT1-MMP [15], MT5-MMP [16], MMP-9, MMP-12 [17], and Presenilin 1 [18]. The induction/upregulation of various MMPs (e.g. MMP-2, MMP-3, MMP-7 and/or MMP-9) has been detected in tumorous liver tissues obtained from, e.g. hepatocellular carcinoma (HCC) patients, whereas the expression of plasminogen activators appears to be largely confined to stromal and inflammatory cells. Inflammation also enhances the expression of MMPs, e.g. in endothelial cells, lymphocytes and macrophages. Furthermore, the expression of certain MMPs and plasminogen activators is also increased in liver cells during liver regeneration [19,20].

Elevated sE-Cad was observed in the serum or plasma of patients with gastric carcinoma [21], bladder cancer [22], non-small cell lung cancer [23], melanoma [24], late-stage colorectal carcinoma [5], and hepatocellular carcinoma [25]. However, higher levels of that cadherin have also been found in gastroesophageal reflux disease [26], multiorgan dysfunction [27], endometriosis [28], familial adenomatous polyposis [25], and acute pancreatitis [29]. Higher amounts of the serum sN-CAD have been described in a group of cancerous patients suffering from prostate carcinoma, breast cancer, ovarian

cancer, and gastrointestinal carcinoma, and have been mentioned as being present in diabetes and liver cirrhosis (data not shown) as well [3]. The present study exhibited a higher plasma concentration of sE-CAD in the liver cancer group than the healthy group. This is consistent with the results reported by Sayoma et al. [25].

Moreover, the level of the marker was observed to be lower in the echinococcosis group than the healthy control group. This suggests that sE-cadherin would be a valuable marker of HCC not only because of its association with early recurrence and extrahepatic metastasis [25], but also owing to its ability to distinguish between liver cancer and echinococcosis. In the present study, the plasma sN-CAD concentration in liver cancer was also observed to be lower than that in the healthy group. These results are difficult to explain and require further analysis. Moreover, they seem to be inconsistent with the results reported by Derycke et al. [3]. However, there is no information about the presence of liver carcinoma in the cancerous group examined by them.

The lower levels of the plasmas sN-CAD and sE-CAD in echinococcosis patients than in healthy subjects could be explained by the ability of *Echinococcus granulosus* to evade certain defence mechanisms by such means as modulating the host's immune system and resisting the host's proteolytic enzymes by dealing efficiently with, e.g. MMP-9 [19,30]. Discriminant analysis revealed that the association of two or more markers significantly increased their efficiency. These results seem to encourage other scientists to combine two or more markers, which are not effective enough when used separately.

Conclusions

Echinococcosis affects the concentration of soluble sE-CAD fragments and Gal-3 in plasma. sE-CAD can be considered as a marker for differentiating between liver cancer and echinococcosis, a parasitic liver disease similar in symptoms. Further study is required to confirm these preliminary results.

Author details

¹Department of Bioanalysis and Drugs Analysis, Faculty of Pharmacy, Medical University of Warsaw, 1 Banacha Street, Warsaw PL-02097, Poland. ²Second Department of General Vascular Oncology Surgery, Medical University of Warsaw, 19/28 Stepiańska Street, Warsaw PL-00739, Poland. ³Department of Gastroenterology, Cardiology and Infectious Diseases, Faculty of Basic Medical Sciences, University of Warmia and Mazury, a 30 Warszawska Street, Olsztyn PL-10082, Poland.

Authors' contributions

All authors participated in the design of the study. MPP applied for and obtained the approval from an ethics committee. MPP and PZ participated in collecting clinical data, performed the clinical diagnosis and samples collecting. JG carried out the ELISA determination, the statistical analysis and drafted the manuscript. MPP, PW, PZ and JP performed reviewing and

editing of the manuscript. JP and PW participated in coordination of the experiment. All authors read and approved the final version of manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 20 December 2011 Accepted: 18 February 2012

Published: 18 February 2012

References

- Moro P, Schantz PM: Echinococcosis: a review. *Int J Infect Dis* 2009, **13**:125-133.
- Reiter-Owona I, Grüner B, Frosch M, Hoerauf A, Kern P, Tappe D: Serological confirmatory testing of alveolar and cystic echinococcosis in clinical practice: results of a comparative study with commercialized and in-house assays. *Clin Lab* 2009, **55**:41-48.
- Derycke L, De Wever O, Stove V, Vanhoecke B, Delanghe J, Depypere H: M B: Soluble N-cadherin in human biological fluids. *Int J Cancer* 2006, **119**:2895-2900.
- Lurisci I, Tinari N, Natoli C, Angelucci D, Cianchetti EaS: Concentrations of Galectin-3 in the Sera of Normal Controls and < p > Cancer Patients. *Clin Cancer Res* 2000, **6**:1389-1393.
- Weiss JV, Klein-Scory S, Kübler S, Reinacher-Schick A, Stricker I, Schmiegel W, Schwarte-Waldhoff I: Soluble E-cadherin as a serum biomarker candidate: Elevated levels in patients with late-stage colorectal carcinoma and FAP. *Int J Cancer* 2011, **128**:1384-1392.
- Weigert J, Neumeier M, Wanninger J, Bauer S, Farkas S, Scherer MN, Schnitzbauer A, Schaffler A, Aslanidis C, Scholmerich J, Buechler C: Serum Galectin-3 is elevated in obesity and negatively correlates with glycosylated hemoglobin in type 2 diabetes. *J Clin Endocrinol Metab* 2010, **95**:1404-1411.
- Lin YH, Lin LY, Wu YW, Chien KL, Lee CM, Hsu RB, Chao CL, Wang SS, Hsein YC, Liao LC, et al: The relationship between serum galectin-3 and serum markers of cardiac extracellular matrix turnover in heart failure patients. *Clin Chim Acta* 2009, **409**:96-99.
- de Boer RaaYL, van Veldhuisen DJ: Galectin-3 in Cardiac Remodeling and Heart Failure. *Curr Heart Fail Rep* 2010, **7**:1-8.
- Matsuda Y, Yamagiwa Y, Fukushima K, Ueno Y, Shimosegawa T: Expression of galectin-3 involved in prognosis of patients with hepatocellular carcinoma. *Hepatology* 2008, **38**:1098-1111.
- Noë V, Fingleton B, Jacobs K, Crawford H, Vermeulen S, Steelant W, Bruyneel E, Matrisian LM, Mareel M: Release of an invasion promoter E-cadherin fragment by matrilysin and stromelysin-1. *J Cell Sci* 2001, **114**:111-118.
- Hayashido Y, Hamana T, Yoshioka Y, Kitano H, Koizumi K, Okamoto T: Plasminogen activator/plasmin system suppresses cell-cell adhesion of oral squamous cell carcinoma cells via proteolysis of E-cadherin. *Int J Oncol* 2005, **27**:693-698.
- Maretzky T, Reiss K, Ludwig A, Buchholz J, Scholz F, Proksch E, de Strooper B, Hartmann D, Saftig P: ADAM10 mediates E-cadherin shedding and regulates epithelial cell-cell adhesion, migration, and beta-catenin translocation. *Proc Natl Acad Sci USA* 2005, **102**:9182-9187.
- Najy AJ, Day KC, Day ML: The ectodomain shedding of E-cadherin by ADAM15 supports ErbB receptor activation. *J Biol Chem* 2008, **283**:18393-18401.
- Reiss K, Maretzky T, Ludwig A, Tousseyn T, de Strooper B, Hartmann D, Saftig P: ADAM10 cleavage of N-cadherin and regulation of cell-cell adhesion and beta-catenin nuclear signalling. *EMBO J* 2005, **24**:742-752.
- Covington MD, Burghardt RC, Parrish AR: Ischemia-induced cleavage of cadherins in NRK cells requires MT1-MMP (MMP-14). *Am J Physiol Renal Physiol* 2006, **290**:F43-F51.
- Monea S, Jordan BA, Srivastava S, DeSouza S, Ziff EB: Membrane localization of membrane type 5 matrix metalloproteinase by AMPA receptor binding protein and cleavage of cadherins. *J Neurosci* 2006, **26**:2300-2312.
- Dwivedi A, Slater SC, George SJ: MMP-9 and -12 cause N-cadherin shedding and thereby beta-catenin signalling and vascular smooth muscle < p > cell proliferation. *Cardiovasc Res* 2009, **81**:178-186.
- Uemura K, Kihara T, Kuzuya A, Okawa K, Nishimoto T, Ninomiya H, Sugimoto H, Kinoshita A, Shimohama S: Characterization of sequential N-cadherin cleavage by ADAM10 and PS1. *Neurosci Lett* 2006, **402**:278-283.
- Marco M, Baz A, Fernandez C, Gonzalez G, Hellman U, Salinas G, Nieto A: A relevant enzyme in granulomatous reaction, active matrix metalloproteinase-9, found in bovine Echinococcus granulosus hydatid cyst wall and fluid. *Parasitol Res* 2006, **100**:131-139.
- Puxbaum V, Mach L: Proteinases and their inhibitors in liver cancer. *World J Hepatol* 2009, **1**:28-34.
- Chan AOO, Lam SK, Chu KM, Lam CM, Kwok E, Leung SY, Yuen ST, Law SYK, Hui WM, Lai KC, et al: Soluble E-cadherin is a valid prognostic marker in gastric carcinoma. *Gut* 2001, **48**:808-811.
- Shariat SF, Matsumoto K, Casella RaJW, Lerner SP: Urinary levels of soluble E-Cadherin in the detection of transitional cell carcinoma of the urinary bladder. *Eur Urol* 2005, **48**:69-76.
- Charalabopoulos K, Gogali A, Dalavaga Y, Daskalopoulos G, Vassiliou M, Bablekos G, Karakosta A, Constantopoulos S: The clinical significance of soluble E-cadherin in nonsmall cell lung cancer. *Exp Oncol* 2006, **28**:83-85.
- Billion K, Ibrahim H, Mauch C, Niessen CM: Increased soluble E-cadherin in melanoma patients. *Skin Pharmacol Physiol* 2006, **19**:65-70.
- Soyama A, Eguchi S, Takatsuki M, Kawashita Y, Hidaka M, Tokai H, Nagayoshi S, Mochizuki S, Matsumoto S, Hamasaki K, et al: Significance of the serum level of soluble E-cadherin in patients with HCC. *Hepatogastroenterology* 2008, **55**:1390-1393.
- Jovov B, Que J, Tobey NA, Djukic Z, Hogan BL, Orlando RC: Role of E-cadherin in the Pathogenesis of Gastroesophageal Reflux Disease. *Am J Gastroenterol* 2011, **106**:1039-1047.
- Pittard AJ, Banks RE, Galley HF, Webster NR: Soluble E-cadherin concentrations in patients with systemic inflammatory response syndrome and multiorgan dysfunction syndrome. *Br J Anaesth* 1996, **76**:629-631.
- Fu C, Lang J: Serum soluble E-cadherin level in patients with endometriosis. *Chin Med Sci J* 2002, **17**:121-123.
- Sewpaul A, French JJ, Khoo TK, Kernohan M, Kirby JA, Charnley RM: Soluble E-cadherin: an early marker of severity in acute pancreatitis. *Book Soluble E-cadherin: an early marker of severity in acute pancreatitis* City; 2009, (Editor ed. ^eds.).
- Monteiro KM, de Carvalho MO, Zaha A, Ferreira HB: Proteomic analysis of the Echinococcus granulosus metacestode during infection of its intermediate host. *Proteomics* 2010, **10**:1985-1999.

doi:10.1186/1746-1596-7-17

Cite this article as: Giebultowicz et al.: How echinococcosis affects potential cancer markers in plasma: galectin-3, sN-cadherin and sE-cadherin? A preliminary report. *Diagnostic Pathology* 2012 **7**:17.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

